

**CUMULATIVE SCIENTIFIC REPORT FOR THE PROJECT  
PN-II-RU-PD-2012-3-0656  
DURING MAY 2013– OCTOBER 2015**

DUMITRU I. STAMATE

GENERAL DATA

**Project** PN-II-RU-PD-2012-3-0656

**Contract no.** 7 / 24.04.2013

**Title:** Combinatorial and homological methods in the study of algebras

**Web page:** <https://dl.dropboxusercontent.com/u/112281424/grantPD-2012-3-0656/index.html>

**Principal Investigator:** dr. Dumitru Ioan Stamate

**Mentor:** C.S.I. dr. Mihai Cipu

**Financing agency:** UEFISCDI

**Host institution:** Institutul de Matematică "Simion Stoilow" al Academiei Române (IMAR), București

**Duration of the project** (updated February 2014): May 2013 – October 2015

**Period covered by this report:** May 2013 – October 2015

CONTENTS

General data	1
1. Brief introduction in the theme of the project	1
2. Scientific objectives	2
3. Main results	3
4. Dissemination of results and other activities	7
References	11

1. BRIEF INTRODUCTION IN THE THEME OF THE PROJECT

Given a graded algebra  $R = \bigoplus_i R_i$  and  $M$  a finitely generated graded  $R$ -module, we are interested in studying the minimal free graded resolution of  $M$  over  $R$ . Its invariants (the Betti numbers  $\beta_i^R(M)$ ) are very important in understanding the equations describing  $M$ . In many relevant cases, e.g.  $R$  is a polynomial ring algebra and  $M = I$  is a graded ideal, there is some extra structure available for this module, and that influences the properties of the module. Several techniques from combinatorics, topology or homological algebra become available. This is an area of much current interest, for details we refer to the monographs [4], [22], [31], [23].

When  $R$  or  $M$  are not already graded, it is a useful way to reduce to the graded case by considering the filtration induced by the powers of an ideal  $\mathfrak{m} \subset R$  and one constructs the associated graded ring

$$\mathrm{gr}_{\mathfrak{m}} R = R/\mathfrak{m} \oplus \mathfrak{m}/\mathfrak{m}^2 \oplus \mathfrak{m}^2/\mathfrak{m}^3 \oplus \cdots$$

and similarly for  $\mathrm{gr}_{\mathfrak{m}} M$ . Of course, several properties may disappear in this transformation, however one knows that the Betti numbers may only increase:  $\beta_i^R(M) \leq \beta_i^{\mathrm{gr}_{\mathfrak{m}} R}(\mathrm{gr}_{\mathfrak{m}} M)$ . We refer to [1] for more details on this topic. From the geometric point of view, if  $(R, \mathfrak{m})$  is the local ring at a point (say the origin) of a variety, its associated graded ring is the ring of coordinates of the tangent cone at the point on the variety, cf. [9], [27].

A special class of graded rings  $R = \bigoplus_i R_i$  is the one where  $R_0$  is a field (or a semisimple ring) and it has an  $R$ -linear resolution. In this case,  $R$  is called Koszul. For an algebra that is not graded *ab initio*, we may use the idea above to pass to the associated graded ring (with respect to a suitable filtration) and we may study the Koszul property for this, as well. This idea has been successfully exploited by V. Reiner and the director of the grant in [25], and there are perspectives to extend its applications.

Already for a standard graded algebra it is rather difficult to check the Koszul property (see [12], [24], [6]). Conca, Trung si Valla ([7]), exploring an idea of Herzog et al([17]) showed that if an algebra  $R$  has a so called *Koszul filtration*, then  $R$  is Koszul. This sufficient condition was useful in determining the Koszul property for several classes of algebras, see the survey [6] and the references within.

## 2. SCIENTIFIC OBJECTIVES

We focused on the following problems and we set the following objectives for the period covered in this report:

- (1) Asymptotic properties of toric algebras.  
Objective: 1 accepted paper in an ISI indexed journal.  
Status: The paper Juergen Herzog, Dumitru I. Stamate, *On the defining equations of the tangent cone of a numerical semigroup ring*, Journal of Algebra vol 418 (2014), 8-28. DOI 10.1016/j.jalgebra.2014.07.008. Preprint version available at arXiv:1308.4644 [math.AC].
- (2) The persistence of homological properties to deformations of Gröbner type.  
Objective: 1 paper submitted to an ISI journal.  
Status:
  - (a) The paper Mircea Cimpoeas, Dumitru I. Stamate, *On intersections of complete intersection ideals*, Preprint 11 pp, 2015 is under review at an international ISI journal.
  - (b) The paper Alexandra Seceleanu, Dumitru I. Stamate, *On Sally semigroup rings*, is in preparation.
- (3) A combinatorial and topological study of the intervals in a numerical semigroup  
Objective: 1 paper submitted.

Status: The paper J. Herzog, D.I. Stamate, *Quadratic numerical semigroups and the Koszul property*, 23 pages. Preprint version available at arXiv:1510.00935 [math.AC] is under review at an international ISI journal.

(4) the Koszul property for some classes of algebras

Objective: 1 preprint

Status: The paper D.I. Stamate, *On the Cohen-Macaulay property for quadratic tangent cones*, preprint 2015.<sup>1</sup>

### 3. MAIN RESULTS

The (numerical) objectives above have been reached through the paper [18] that was published by the Journal of Algebra, through the preprints [5] and [19] which are under evaluation at prestigious ISI journals, and through the preprint [30] which is currently circulated among some specialists before being submitted to a journal.

We also mention the preprint [28], realised outside the minimal objectives we had set. The latter preprint will be finalized soon.

In what follows we briefly describe the contents of the papers already written in this project.

Our paper [18] has appeared:

Jürgen Herzog, **Dumitru I. Stamate**

*On the defining equations of the tangent cone of a numerical semigroup ring*, J. Algebra **418** (2014), 8–28.

We describe its contents, that will be useful to describe the other papers, as well.

Let  $\mathbf{a} = a_1 < \dots < a_r$  be a sequence of nonnegative integers. We denote  $\langle a_1, \dots, a_r \rangle$  (or simply  $\langle \mathbf{a} \rangle$ ) the subsemigroup of  $\mathbb{N}$  generated by  $a_1, \dots, a_r$ . In other words,  $\langle \mathbf{a} \rangle$  consists of all  $\mathbb{N}$ -linear combinations of  $a_1, \dots, a_r$ . If  $H = \langle a_1, \dots, a_r \rangle$  we call  $a_1, \dots, a_r$  a *generating system* for  $H$ . From now on, any subsemigroup  $H \subset \mathbb{N}$  with  $0 \in H$  will be called a *numerical semigroup*. Such a semigroup is finitely generated and it admits a unique minimal generating system whose cardinality we denote by  $\mu(H)$ . It is usually part of the definition of numerical semigroups that the greatest common divisor of its generators be 1. In the context of this paper it is convenient to drop this requirement.

For any integer  $k$ , let  $\mathbf{a} + k$  denote the shifted sequence  $a_1 + k, \dots, a_r + k$ . If  $H$  is minimally generated by  $\mathbf{a} = a_1, \dots, a_r$ , we let  $H_k = \langle \mathbf{a} + k \rangle$ . We refer to the family of semigroups  $\{H_k\}_{k \in \mathbb{N}}$  as the *shifted family* associated to  $H$ . One can see that although the  $a_i$ 's generate  $H$  minimally, for some shifts  $k$  the sequence  $\mathbf{a} + k$  may not be a minimal generating system for  $H_k$ . In particular,  $(H_k)_\ell$  may differ from  $H_{k+\ell}$ . For example, when  $H = \langle 3, 5, 7 \rangle$  we get  $H_1 = \langle 4, 6, 8 \rangle = \langle 4, 6 \rangle$ . On the other hand, if  $H = \langle \mathbf{a} \rangle$  is minimally generated by  $\mathbf{a} = a_1 < \dots < a_r$ , then for all  $k > a_r - 2a_1$ ,  $H_k$  is minimally generated by the sequence  $\mathbf{a} + k$ .

Let  $K$  be any field and  $S = K[x_1, \dots, x_r]$  the polynomial ring over  $K$  in the variables  $x_1, \dots, x_r$ . Let  $\mathbf{a} = a_1 < \dots < a_r$  be a sequence of positive integers and

---

<sup>1</sup>Updated Dec 2015: 20 pages. Preprint version available at arXiv:1512.04893 [math.AC]. Submitted.

$\varphi : S \rightarrow K[t]$  is the  $K$ -algebra map letting  $\varphi(x_i) = t^{a_i}$  for  $i = 1, \dots, r$ , where  $K[t]$  is the polynomial ring over  $K$  in the variable  $t$ . If we denote  $H = \langle a_1, \dots, a_r \rangle$ , then the image of  $\varphi$  is the semigroup ring  $K[H]$ , i.e. the  $K$ -subalgebra of  $K[t]$  generated by  $t^{a_1}, \dots, t^{a_r}$  over  $K$ . Let  $I(\mathbf{a})$  be the kernel of  $\varphi$ . When  $\mathbf{a}$  is a minimal generating system for  $H$ , the ideal  $I(\mathbf{a})$  depends only on  $H$  and we denote  $I_H = I(\mathbf{a})$ .

One knows from [16] that the minimal number of generators  $\mu(I_H)$  of  $I_H$  is at most 3 for  $r \leq 3$ . On the other hand, already for  $r = 4$ , the value of  $\mu(I_H)$  may be arbitrarily large, cf. [3]. The more it is surprising that for any numerical semigroup  $H$  there exists an upper bound for  $\mu(I_{H_k})$  independent of  $k$ , see [32, Theorem 1.1]. This result was conjectured by J. Herzog and H. Srinivasan and it was proved first by P. Gimenez, I. Sengupta și H. Srinivasan în [14] for numerical semigroups generated by an arithmetic sequence. This conjecture and a stronger form of it have been recently proved by T. Vu in [32].

Although for  $r = 3$  we have a  $\mu(I_H) \leq 3$ , however the number of generators for  $I_H^*$  may be arbitrarily large. A first family of such examples was found by T. Shibuta, see [15]. For this family its *width* is not bounded, where by the width of the semigroup  $H$ , denoted  $\text{wd}(H)$ , we understand the difference between the largest and the smallest element in the minimal generating system of  $H$ . In our Corollary 1.16 we prove that there exists un upper bound for  $\mu(I_H^*)$  which is valid for all semigroups of fixed width. This follows from a recent theorem of Vu, cf. [32, Theorem 1.1] and our next theorem.

**Theorem 1.4.** *Let  $H$  be a numerical semigroup. There exists  $k_0 \in \mathbb{N}$  such that for  $k \geq k_0$ , the ideal  $I_{H_k}$  is minimally generated by a standard basis, and  $\beta_i(I_{H_k}) = \beta_i(I_{H_k}^*)$  for all  $i$ . In particular,  $\text{gr}_m K[H_k]$  is Cohen–Macaulay for all  $k \geq k_0$ .*

The methods that we use for showing the existence of a uniform upper bound for  $\mu(I_H^*)$  for all numerical semigroups  $H$  of fixed width do not give an explicit value. However, numerical experiments with SINGULAR [8] give us reasons to believe that  $\binom{\text{wd}(H)+1}{2}$  is such an upper bound, and it can not be improved since it is reached by numerical semigroups generated by certain intervals of integers. We prove that this conjectured bound is valid for any numerical semigroup with the property that  $\mu(I_H^*) \leq \mu(I_{\tilde{H}}^*)$ , where  $\tilde{H}$  is the semigroup generated by all integers in the interval whose ends are the smallest and the largest minimal generator of  $H$ .

To support our conjecture, we prove in Proposition 2.10 that for a numerical semigroup  $H$  generated by an arithmetic sequence we even have  $\beta_i(I_H^*) \leq \beta_i(I_{\tilde{H}}^*)$ , for all  $i$ . It is possible that such an inequality take place for any numerical semigroup!

In the last section of the paper we study several families of semigroups where we tested the above conjectures and we describe the ideal  $I_H^*$  for all members  $H$  in such families. The first family is based on a well-known result of J. Sally in [26], where the author describes the equations of the tangent cone of a Gorenstein local ring with the property  $r = e + d - 3$ . Here  $r$  denotes the embedding dimension,  $e$  denotes the multiplicity, and  $d$  is the Krull dimension of the ring. We define a *Sally semigroup* to be a numerical semigroup whose semigroup ring satisfies the above identity. We prove that there exists Sally semigroups of any multiplicity  $e \geq 4$ . Another family

we study is due to H. Bresinsky [3]. This is the first known family of 4-generated numerical semigroups such that  $\mu(I_H)$  may be arbitrarily large when  $H$  runs in this family. We show that a Bresinsky semigroup is Cohen-Macaulay and we exhibit a minimal generating system of equations that is also a standard basis.

The other two families refer to 3-generated semigroups, and although their members have arbitrary width, the behaviour of  $\mu(I_H^*)$  is quite different from above. For  $a > 3$ , the ideal  $I_H^*$  associated to the semigroup  $H = \langle a, a + 1, 2a + 3 \rangle$  is generated by  $\lfloor \frac{a-1}{3} \rfloor + 3$  monomials. For that family, the number of generators of  $I_H^*$  is a quasi-linear function on the width of  $H$ , and it goes to infinity as  $\text{wd}(H)$  goes to infinity, too. When  $a = 3b$  we come across an example found by T. Shibuta, that was treated in [15, Example 5.5] with different methods.

On the other hand, for any  $a, b > 3$  coprime, if we let  $H = \langle a, b, ab - a - b \rangle$ , then also  $\mu(I_H^*) = 4$ , although the widths of such semigroups may be arbitrarily large.

The work on the paper [5]:

Mircea Cimpoeaş, **Dumitru I. Stamate**,

*On intersections of complete intersection ideals*, 11 pp, preprint 2015. Submitted.

was stimulated by numerical experiments and by the experience gained when preparing the previously described paper.

An ideal  $I$  in a (Noetherian) ring  $R$  is called a *complete intersection* (CI for short) if it can be generated by height  $I$  elements, i.e. the minimum allowed by a famous theorem of Krull. This class of Gorenstein rings has many homological properties and a geometric relevance. Therefore its extremal character has attracted the attention of many researchers.

In general the class of CI ideals is not closed to the main operations with ideals, e.g. sum, intersection, except some very trivial cases. However, in this paper we present infinite families of CI ideals that are (each) closed to taking intersections.

In order to explain the constructions we continue to use the notation introduced in the previous paragraphs. We saw that for a list  $\mathbf{a}$  and a shift  $k \gg 0$ , the Betti numbers for  $I_{\mathbf{a}+k}$  are periodic in  $k$ . In particular, if  $I_{\mathbf{a}+k}$  is CI for some  $k \gg 0$ , then there exist an infinity of shifts  $j$  such that  $I_{\mathbf{a}+j}$  and  $I_{\mathbf{a}+j}^*$  are also CI.

For a finite family of indices  $\mathcal{A} \subset \mathbb{N}$  we consider

$$\mathcal{I}_{\mathcal{A}} = \bigcap_{j \in \mathcal{A}} I_{\mathbf{a}+j}$$

and similarly

$$\mathcal{J}_{\mathcal{A}} = \bigcap_{j \in \mathcal{A}} I_{\mathbf{a}+j}^*.$$

These ideals are no longer binomial and many of the properties of toric ideals are lost. Using Gröbner bases techniques and the work in [20] we prove that when  $I_{\mathbf{a}+j}$  is CI for all  $j \in \mathcal{A}$  and  $\min \mathcal{A} \gg 0$ , then  $\mathcal{I}_{\mathcal{A}}$  and  $\mathcal{J}_{\mathcal{A}}$  are also CI.

More precise results are obtained for 3-generated semigroups using the characterization of shifts  $k \gg 0$  such that  $I(\mathbf{a} + k)$  is CI, cf. [29].

Numerical experiments make us believe that this stability to intersections of CI ideals in a shifted family of semigroups is only a facet of a larger periodicity phenomenon. Also, Gröbner deformations of these shifted ideals have the same periodicity to intersections. We believe these are reasons for the further study of the "asymptotic properties" and/or the Gröbner deformations of the (ideal of) affine semigroup rings, not necessarily numerical.

In [19]:

Juergen Herzog, **Dumitru I. Stamate**, *Quadratic numerical semigroups and the Koszul property*, 23 pages. Preprint version available at arXiv:1510.00935 [math.AC]. Submitted.

we find effective bounds for the multiplicity  $e(H)$  of a numerical semigroup  $H$  such that its associated graded ring  $\text{gr}_m K[H]$  is defined by quadratic relations, or it is even Koszul. We call  $H$  quadratic/Koszul if  $\text{gr}_m K[H]$  is so.

**Theorem 1.1.** ([19]) *Let  $H$  be a quadratic numerical semigroup minimally generated by  $n$  elements, and  $K[H]$  the associated semigroup ring. Then*

- (a)  $n \leq e(H) \leq 2^{n-1}$ ;
- (b)  $e(H) = n \iff I_H^*$  has a linear resolution;
- (c)  $e(H) = 2^{n-1} \iff I_H^*$  is a complete intersection ideal  $\iff I_H$  is a complete intersection ideal.

An interesting problem is to effectively determine all the possible multiplicities in the interval  $[n, 2^{n-1}]$ . We estimate that not all these values are possible. In this direction we have the following result:

**Theorem 1.9.** ([19]) *Let  $H$  be a quadratic numerical semigroup minimally generated by  $n$  elements such that  $\text{gr}_m K[H]$  is Cohen-Macaulay. Then*

- (a) either  $n \leq e(H) \leq 2^{n-1} - 2^{n-3}$ , or  $e(H) = 2^{n-1}$ ;
- (b) If  $e(H) = 2^{n-1} - 2^{n-3}$ , then  $I_H^*$  is an almost complete intersection ideal.

In case of (b), the ideal  $I_H^*$  has a quadratic Gröbner basis with respect to revlex induced by  $x_n > \dots > x_1$ .

One knows that in general, if the ideal  $I$  in the polynomial ring  $S$  is  $G$ -quadratic (i.e. after an eventual linear change of coordinates it admits a quadratic Gröbner basis with respect to a monomial term order), then  $S/I$  is Koszul. We prove that:

**Proposition 1.12.** ([19]) *If  $H$  is a quadratic numerical semigroup with  $\text{emb dim}(H) = n$ , and  $e(H)$  equals  $n$ ,  $2^{n-1}$  or also  $2^{n-1} - 2^{n-3}$  in case  $\text{gr}_m K[H]$  is Cohen-Macaulay, then  $H$  is  $G$ -quadratic.*

New numerical semigroups can be obtained by gluing two semigroups with fewer generators. If  $H_1$  and  $H_2$  are two numerical semigroups,  $c_1$  and  $c_2$  are coprime integers such that  $c_1 \in H_2 \setminus G(H_2)$  and  $c_2 \in H_1 \setminus G(H_1)$ , we say that  $H = \langle c_1 H_1, c_2 H_2 \rangle$  is obtained by gluing  $H_1$  and  $H_2$ . The most famous result in this direction is Delorme's characterisation of complete intersection (CI) numerical semigroups. More precisely, any such semigroup is obtained via a series of gluings starting with  $\mathbb{N}$ .

If  $c_1 = 2$  and  $H_2 = \mathbb{N}$  we say that  $H = \langle 2H_1, c_2 \rangle$  is obtained from  $H_1$  by a quadratic gluing. An important result obtained by us (Theorem 2.14, [19]) is that any quadratic CI numerical semigroup can be obtained by a series of quadratic

gluings starting from  $\mathbb{N}$ . This is a consequence of the fact that the semigroup  $H = \langle 2L, \ell \rangle$  is quadratic/Koszul/ $G$ -quadratic if and only if  $L$  has the respective property.

In the last section of the paper we apply the previous results to completely describe the quadratic semigroups among those generated by arithmetic or geometric sequences, or 3-generated, and also symmetric or pseudo-symmetric 4-generated.

In [30]:

**Dumitru I. Stamate**, *On the Cohen-Macaulay property for quadratic tangent cones*, preprint 2015.

we continue the line of research initiated in [19]. We show that any 4-generated numerical semigroup, if it is quadratic, then it is also  $G$ -quadratic, and moreover  $\text{gr}_{\mathfrak{m}} K[H]$  is even Cohen-Macaulay. In case  $\text{emb dim}(H) = 5$ , as a result of a detailed analysis, we show that if  $H$  is quadratic, then  $\text{gr}_{\mathfrak{m}} K[H]$  is Cohen-Macaulay except two families of semigroups that are explicitly described.

These results allow us to obtain the effective multiplicities such that  $H$  is quadratic, and  $\text{emb dim}(H) \leq 5$ . Moreover, we can determine the  $h$ -vector of  $\text{gr}_{\mathfrak{m}} K[H]$ , and as an application we have that under the above conditions, if the field  $K$  is algebraically closed and of characteristic  $\neq 2$  and  $H$  is Koszul, then  $H$  is also  $G$ -quadratic.

For any  $n > 5$  we construct infinitely many examples of semigroups  $H$  that are Koszul, even  $G$ -quadratic, and such that  $\text{gr}_{\mathfrak{m}} K[H]$  is not Cohen-Macaulay. This result is important, since random computer search did not offer us any such example.

In [28]:

Alexandra Seceleanu, **Dumitru I. Stamate**, *On Sally semigroup rings*, in preparation.

we study the class of Sally semigroups described above. We have experimentally noticed that for artinian Sally rings, their Betti numbers are the same. We confirm this observation and we give a precise formula for them. We employ recent results of Elias and Rossi ([10]) about the classification of short artinian algebras, and also a theorem regarding the decomposition into connected sums of rings, cf. Ananthnarayan et al ([2]). Such a formula is useful because for the Sally numerical semigroup we presented in [18] their Betti numbers are rather large, close to the maximum conjectured in [18].

Complementing [18], we present many new examples of Sally numerical semigroups.

#### 4. DISSEMINATION OF RESULTS AND OTHER ACTIVITIES

Mobilities played an important part in the planning of the activities. We considered to be very important to meet with experts in our area of research from prestigious universities. On one hand we could present our results, and on the other hand these visits materialized into several research projects initiated within the theme of our project, see the previous section.

I was invited to give seminar talks at all the universities that I visited. These presentations have been excellent opportunities to obtain feed-back and suggestions for subsequent developments. Next we outline the research visits we made, which in general goes hand in hand with the dissemination talks.

**Research visits:**

- (1) University Osnabrueck, Germania, June 2013. Host: prof. Tim Römer.
- (2) University Duisburg-Essen, Essen, Germania, July – August 2013. Host: prof. Jürgen Herzog.
- (3) University of Nebraska, Lincoln, NE, SUA, September 2013. Host: prof. Roger Wiegand.
- (4) University of Minnesota, Minneapolis, MN, SUA, October 2013. Host: prof. Victor Reiner.
- (5) University of Missouri, Columbia, MO, SUA, October 2013. Host: prof. Hema Srinivasan.
- (6) University of Nebraska, Lincoln, NE, SUA, January 2014. Host: prof. Roger Wiegand.
- (7) University of Minnesota, Minneapolis, MN, SUA, February 2014. Host: prof. Victor Reiner.
- (8) Universite de Montpellier 2, Institut de Mathematiques et de Modelisations de Montpellier, Montpellier, France, May 2014. Host: dr. Ignacio Garcia Marco.
- (9) Universita di Genova, Italy, June 2014. Host: prof. Aldo Conca.
- (10) University Duisburg-Essen, Essen, Germania, July – August 2014. Host: prof. Jürgen Herzog.
- (11) University of Nebraska, Lincoln, NE, SUA, September 2014. Host: dr. Alexandra Seceleanu.
- (12) University of Sheffield, UK, May 2015 (1 week). Hosts: prof. Moty Katzman and Dr. Ines Henriques.
- (13) University Duisburg-Essen, Essen, Germany, Septembrie 2015 (1 week). Host: prof. Jürgen Herzog.

**Invited experts:**

During 2–7 September 2014, prof. Jorge Ramirez Alfonsin, Université Montpellier 2, France visited us. On this occasion he gave a series of 4 talks about the algebraic properties of matroids, in the framework of the National Algebra School.

We also discussed his very recent results concerning the Moebius function of intervals in numerical semigroups and new combinatorial approaches in the study of (numerical) semigroups.

During 12–20 June 2015, prof. Juergen Herzog from Universitaet Duisburg-Essen visited us, on which occasion we discussed about our ongoing projects, including [19]. On 16 June 2015 he gave a very well attended lecture at IMAR on *Ideals associated to isotone maps between finite posets*.



## Dissemination of results:

- (1) D. Stamate, *On the CI property of the tangent cone of a toric ring*, Workshop for Young Researchers in Mathematics, Ovidius University Constanța, 8–10 May 2013.
- (2) D. Stamate, *Shifting semigroups*, short talk, Workshop "Syzygies in Berlin", Freie Universität, Berlin, Germania, 28 May 2013.
- (3) D. Stamate, *Shifted semigroup rings*, Oberseminar University of Osnabrueck, Germania, 4 June 2013.
- (4) D. Stamate, *On the CI property of the tangent cone of a toric ring*, AMS-RMS Joint meeting, Special Session on Commutative Algebra, Alba Iulia, 30 June 2013.
- (5) D. Stamate, *On the equations of toric rings*, University Duisburg-Essen, Essen, 29 August 2013.
- (6) D. Stamate, *Tools of Combinatorial Commutative Algebra 2*, National Algebra School "Algebraic methods in Combinatorics", 3 September 2013.
- (7) D. Stamate, *Matroids and realisability*, National Algebra School "Algebraic methods in Combinatorics", 4 September 2013.
- (8) D. Stamate, *On the defining equations of the tangent cone of a numerical semigroup ring*, Comm. Algebra Seminar talk, University of Nebraska, Lincoln, NE, SUA, 18 September 2013.
- (9) D. Stamate, *On the defining equations of the tangent cone of a numerical semigroup ring*, Comm. Algebra Seminar talk, University of Minnesota, Minneapolis, MN, SUA, 14 October 2013.
- (10) D. Stamate, *On the defining equations of the tangent cone of a numerical semigroup ring*, Comm. Algebra Seminar talk, University of Missouri, Columbia, MO, SUA, 22 October 2013.
- (11) D. Stamate, *Asymptotic properties of numerical semigroups I, II*, Commutative Algebra seminar IMAR & Univ. București, 19 și 26 November 2013.
- (12) D. Stamate, *About the structure of Sally rings*, Comm. Algebra Seminar talk, University of Minnesota, Minneapolis, MN, SUA, 14 February 2014.
- (13) D. Stamate, *On the CI property of the tangent cone of a toric ring*, Workshop for Young Researchers in Mathematics, Ovidius University Constanța, 22–23 May 2014.
- (14) D. Stamate, *On numerical semigroup rings and their defining relations*, Séminaire Algèbre et géométrie combinatoires, Université de Montpellier 2, Franța, 27 May 2014.
- (15) D. Stamate, *On the defining equations of the tangent cone of a numerical semigroup ring*, Commutative Algebra Seminar talk, University of Genova, Italia, 3 June 2014.
- (16) D. Stamate, *Asymptotic properties of numerical semigroups*, National Algebra School "Algebraic and Combinatorial Applications of Toric Ideals", 3 September 2014.
- (17) D. Stamate, *Flavors of Koszul rings*, Comm. Algebra Seminar talk, University of Nebraska, Lincoln, NE, SUA, 17 September 2014.

- (18) D. Stamate, *On intersections of complete intersection ideals*, Comm. Algebra Seminar IMAR and Univ. Bucharest, 24 February 2015.
- (19) D. Stamate, *Koszul filtrations*, Comm. Algebra Seminar IMAR and Univ. Bucuresti, 28 April 2015.
- (20) D. Stamate, *Koszul filtrations (II). Grobner flags.*, Comm. Algebra Seminar IMAR and Univ. Bucharest, 5 May 2015.
- (21) D. Stamate, *Koszul filtrations (III). Strongly Koszul rings and the ungraded version.*, Comm. Algebra Seminar IMAR and Univ. Bucharest, 12 May 2015.
- (22) D. Stamate, *Koszul rings and the combinatorics of posets.*, Comm. Algebra Seminar IMAR and Univ. Bucharest, 19 May 2015.
- (23) D. Stamate, *Filtrations for Koszul rings*, Workshop for Young Researchers in Mathematics, Universitatea Ovidius Constanta, Romania, 21 May 2015.
- (24) D. Stamate, *On the defining equations of tangent cones of numerical semigroup rings*, Algebra / Algebraic Geometry seminar, University of Sheffield, United Kingdom, 27 May 2015.
- (25) D. Stamate, *Ungraded strongly Koszul rings*, The Eighth Congress of Romanian Mathematicians, Iasi, Romania, 26 June-1 July 2015.
- (26) D. Stamate, *On the Koszul property for numerical semigroup rings*, Syzygies in Algebra and Geometry 2015, Busan, Korea, 26-30 August 2015.
- (27) D. Stamate, *Koszul filtrations for ungraded rings*, National Algebra School – Interactions of Computer Algebra with Commutative Algebra, Combinatorics and Algebraic Statistics, IMAR, Bucharest, 3 Septembrie 2015.
- (28) D. Stamate, *On the Koszul property for numerical semigroup rings (I, II)*, Comm. Algebra Seminar IMAR and Univ. Bucharest, 6 and 13 October 2015.
- (29) D. Stamate, *Quadratic numerical semigroups and the Koszul property*, The first Romanian-Turkish Mathematics Colloquium, Constanta, Romania, 16 October 2015.
- (30) D. Stamate, *The structure of quadratic CI semigroups*, Comm. Algebra Seminar IMAR and Univ. Bucharest, 10 November 2015.

**Other seminar talks:**

- (1) M. Cipu, *Algebraic tools for discrete tomography*, Commutative Algebra seminar IMAR & Univ. București, 18 February 2014.
- (2) M. Cipu, *Quantic Ehrhart polynomials*, Commutative Algebra seminar IMAR & Univ. București, 1 April 2014.
- (3) D. Stamate, *On the subadditivity problem for maximal shifts in free resolutions, after Herzog et al*, Commutative Algebra seminar IMAR & Univ. București, 6 May 2014.
- (4) M. Cipu, *A conjectural characterization for complete intersection numerical semigroups*, Commutative Algebra seminar IMAR & Univ. București, 4 November 2014.

**Other visits** supported by this project:

- (1) Conference GMZ50 honoring Gunter Ziegler, Freie Universität Berlin, Germania, organized by Christian Haase, Raman Sanyal, Nadja Wisniewski, 25 May 2013.
- (2) COCOA School, Universität Osnbrück, Germania, organized by W. Bruns, L. Robbiano, A. Bigatti, 10–14 June 2013.
- (3) ETMAGT-International Conference Experimental and Theoretical Methods in Algebra, Geometry and Topology, Eforie Nord, 20–24 June 2013. (Dumitru Stamate and Mihai Cipu)
- (4) Recent Trends in Algebraic and Geometric Combinatorics, Madrid, 26-30 November 2013. (Mihai Cipu)
- (5) Encuentros de Algebra Computacional y Aplicaciones-EACA, Barcelona, Spania, 17–22 June 2014. (Mihai Cipu)
- (6) Meeting On Combinatorial Commutative Algebra, MOCCA, Levico Terme, Italia, 8–12 September 2014. (Mihai Cipu)

The discussions with prof. Tim Römer on the occasion of the visit to Osnbrück in June 2013 favored the **organization** of the 21st edition of the National Algebra School – ”Algebraic methods in Combinatorics” where prof. Tim. Römer was a keynote speaker. We organized this event at IMAR, 2-6 September 2013, together with Viviana Ene, Mihai Epure, Miruna Roşca, Andrei Zarojanu. Scientific committee: Dorin Popescu, Tim Römer, Marius Vlădoiu.

In 2014, on the occasion of the visit to Montpellier I invited prof. Jorge Ramirez-Alfonsin to give a series of lectures in Bucharest at the 22nd edition of the National Algebra School–”Algebraic and Combinatorial Applications of Toric Ideals”. Scientific committee: Hara Charalambous, Mihai Cipu (member of the team of the grant), Jorge Ramirez Alfonsin. We organized this school at IMAR, 1-5 September 2014, together with Florin Ambro, Viviana Ene, Mihai Epure, Miruna Roşca, Marius Vlădoiu and Andrei Zarojanu.

In 2015 I co-organized the 23rd edition of the National Algebra School - Interactions of Computer Algebra with Commutative Algebra, Combinatorics and Algebraic Statistics, 31 August - 4 September 2015, IMAR, Bucharest, together with Viviana Ene, Mihai Epure, and Marius Vlădoiu. the scientific committee: Gerhard Pfister, Dorin Popescu.

At all three editions there were many participants, including students and PhD students.

On the occasion of the 8th Congress of Romanian Mathematicians in Iasi, Romania, together with Dorin Popescu, I co-organized the Special Session ”Local rings and homological algebra, dedicated to Prof. Nicolae Radu” , 27 June 2015.

## REFERENCES

- [1] R. Achilles, L. L. Avramov, *Relations between properties of a ring and of its associated graded ring*, Seminar D. Eisenbud/B. Singh/W. Vogel, Vol. 2, Teubner, Leipzig, 1982, 5–29.
- [2] H. Ananthnarayan, Ela Celikbas, Zheng Yang, *Decomposing Gorenstein Rings as Connected Sums*, preprint, arXiv:1406.7600 [math.AC].

- [3] H. Bresinsky, *On prime ideals with generic zero  $x_i = t^{n_i}$* , Proc. Amer. Math. Soc. **47** no.2 (1975), 329–332.
- [4] W. Bruns, J. Herzog, *Cohen–Macaulay Rings*, Cambridge University Press, 2nd edition, 1998.
- [5] M. Cimpoeaş, D.I. Stamate, *On intersections of complete intersection ideals*, 11 pp, preprint 2015. Submitted.
- [6] A. Conca, E. De Negri, M.E. Rossi, *Koszul algebras and regularity*. Commutative algebra, 285–315, Springer, New York, 2013.
- [7] A. Conca, N.V. Trung, G. Valla, *Koszul property for points in projective spaces*. Math. Scand. **89** (2001) no. 2, 201–216.
- [8] W. Decker, G.-M. Greuel, G. Pfister, H. Schönemann, SINGULAR 3-1-6 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2012).
- [9] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics **150**, Springer, 1995.
- [10] J. Elias, M.E. Rossi, *Isomorphism classes of short Gorenstein local rings via Macaulays inverse system*, Trans. Amer. Math. Soc., **364** (2012), 4589–4604.
- [11] V. Ene, J. Herzog, *Gröbner bases in commutative algebra*, Graduate Studies in Mathematics **130**, American Mathematical Society, 2012.
- [12] R. Fröberg, *Koszul algebras* in Advances in commutative ring theory (Fez, 1997), 337–350, Lecture Notes in Pure and Appl. Math., 205, Dekker, New York, 1999.
- [13] P. Gimenez, I. Sengupta, H. Srinivasan, *Minimal free resolutions for certain affine monomial curves*, in A. Corso, C. Polini (Eds.), Commutative Algebra and Its Connections to Geometry, PASI 2009, in: Contemp. Math., vol **555**, Amer. Math. Soc., 2011, pp. 87–95.
- [14] P. Gimenez, I. Sengupta, H. Srinivasan, *Minimal graded free resolutions for monomial curves defined by arithmetic sequences*, J. Algebra **338** (2013), 294–310.
- [15] S. Goto, W. Heinzer, M. Kim, *The leading ideal of a complete intersection of height two. II*, J. Algebra **312** (2007), 709–732.
- [16] J. Herzog, *Generators and relations of Abelian semigroups and semigroup rings*, Manuscripta Math. **3** (1970), 175–193.
- [17] J. Herzog, T. Hibi, G. Restuccia, *Strongly Koszul algebras*, Math. Scand. **86** (2000), 161–178.
- [18] J. Herzog, D.I. Stamate, *On the defining equations of the tangent cone of a numerical semigroup ring*, J. Algebra **418** (2014), 8–28. arXiv:1308.4644 [math.AC].
- [19] J. Herzog, D.I. Stamate, *Quadratic numerical semigroups and the Koszul property*, preprint 23 pp, 2015, arXiv:1510.00935 [math.AC]. Submitted.
- [20] A. V. Jayanthan, H. Srinivasan, *Periodic Occurance of Complete Intersection Monomial Curves*, Proc. AMS, **141** (2013), 4199–4208. arXiv:1203.1991 [math.AC].
- [21] *MAGMA-Computational Algebra System*, developed by Computational Algebra Group at the University of Sydney, Australia, available at <http://magma.maths.usyd.edu.au>.
- [22] E. Miller, B. Sturmfels, *Combinatorial Commutative Algebra*, Springer GTM, 2005.
- [23] I. Peeva, textitGraded Syzygies, Algebra and Applications Ser., Vol 14, Springer 2011.
- [24] A. Polishchuk, L. Positselski, *Quadratic algebras*, University Lecture Series, 37, American Mathematical Society, Providence, RI, 2005.
- [25] V. Reiner, D. I. Stamate, *Koszul incidence algebras, affine semigroup rings, and Stanley-Reisner ideals*, Adv. Math. **224**, no. 6 (2010), 2312–2345.
- [26] J.D. Sally, *Good embedding dimensions for Gorenstein singularities*, Math. Ann. **249** (1980), 95–106.
- [27] I. R. Şafarevici, *Bazele geometriei algebrice*, Ed. St. Encicl., Bucureşti, 1976.
- [28] A. Seceleanu, D.I. Stamate, *On Sally semigroup rings*, in preparation.
- [29] D. I. Stamate, *Asymptotic properties in the shifted family of a numerical semigroup with few generators*, Semigroup Forum, in press. available online, DOI 10.1007/s00233-015-9724-2.
- [30] D. I. Stamate, *On the Cohen-Macaulay property for quadratic tangent cones*, preprint 2015.
- [31] R. P. Stanley, *Combinatorics and Commutative Algebra*, second edition, Birkäuser Boston, 1996.

[32] T. Vu, *Periodicity of Betti numbers of monomial curves*, J. Algebra **418** (2014), 66–90.  
arXiv:1304.1659 [math.AC].

30 October 2015  
dr. Dumitru I. Stamate

DUMITRU I. STAMATE, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY  
OF BUCHAREST, STR. ACADEMIEI 14, BUCHAREST, ROMANIA, AND

SIMION STOILLOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, RESEARCH  
GROUP OF THE PROJECT PN-II-RU-PD-2012-3-0656, P.O.Box 1-764, BUCHAREST 014700,  
ROMANIA

*E-mail address:* `dumitru.stamate@fmi.unibuc.ro`